

原著論文

- 1) Total biosynthesis of opiates by stepwise fermentation using engineered *Escherichia coli*. Nakagawa A., Matsumura E., Koyanagi T., Katayama T., Kawano N., Yoshimatsu K., Yamamoto K., Kumagai H., Sato F. and Minami H. *Nature Communications* 7, 10390 (2016).
- 2) Tracing microbiota changes in yamahai-moto, the traditional Japanese sake starter. Koyanagi T., Nakagawa A., Kiyohara M., Matsui H., Tsuji A., Barla F., Take H., Katsuyama Y., Tokuda K., Nakamura S., Minami H., Enomoto T., Katayama T. and Kumagai H. *Bioscience, Biotechnology, and Biochemistry*. 80, 399-406 (2016).
- 3) α -Amylase from Mon Thong durian (*Durio zibethinus* Murr. cv. Mon Thong)-nucleotide sequence analysis, cloning and expression. Posoongnoen S., Ubonbal R., Thammasirirak S., Daduang J., Minami H., Yamamoto K., Daduang S. *Plant Biotechnology* 32, 1-10 (2015).
- 4) (R,S)-Tetrahydropapaveroline production by stepwise fermentation using engineered *Escherichia coli*. Nakagawa A., Matsuzaki C., Matsumura E., Koyanagi T., Katayama T., Yamamoto K., Sato F., Kumagai H. and Minami H. *Scientific Reports* 4, 6695 (2014).
- 5) Asymmetric synthesis of tetrahydroisoquinolines by enzymatic Pictet-Spengler reaction. Nishihachijo M., Hirai Y., Kawano S., Nishiyama A., Minami H., Katayama T., Yasohara Y., Sato F., Kumagai H. *Bioscience, Biotechnology, and Biochemistry* 78, 701-707 (2014).
- 6) Improvement of reticuline productivity from dopamine by using engineered *Escherichia coli*. Kim J.S., Nakagawa A., Yamazaki Y., Matsumura E., Koyanagi T., Minami H., Katayama T., Sato F. and Kumagai H. *Bioscience, Biotechnology, and Biochemistry* 77, 2166-2168 (2013).
- 7) Pyrosequencing analysis of microbiota in Kaburazushi, a traditional medieval sushi in Japan. Koyanagi T., Nakagawa A., Kiyohara M., Matsui H., Yamamoto K., Barla F., Take H., Katsuyama Y., Tsuji A., Shijimaya M., Nakamura S., Minami H., Enomoto T., Katayama T., Kumagai H. *Bioscience, Biotechnology, and Biochemistry* 77, 2125-2130 (2013).
- 8) Pathway engineering of benzyloquinoline alkaloid biosynthesis in transgenic California poppy cells with ectopic expression of tetrahydroberberine oxidase from *Coptis japonica*. Matsushima Y., Minami H., Hori K. and Sato F. *Plant Biotechnology* 29, 473-481 (2012).
- 9) Eukaryotic-type aromatic amino acid decarboxylase from root colonizer *Pseudomonas putida* is highly specific for 3,4-dihydroxyphenyl-L-alanine (DOPA), an allelochemical in soil. Koyanagi T., Nakagawa A., Sakurama H., Yamamoto K., Sakurai N., Takagi T., Minami H., Katayama T. and Kumagai H. *Microbiology* 158, 2965-2974 (2012).
- 10) Bench-top fermentative production of plant benzyloquinoline alkaloids using a bacterial platform. Nakagawa A., Minami H., Kim JS., Koyanagi T., Katayama T., Sato F. and

- Kumagai H. *Bioengineered Bugs* 3, 49-53 (2012).
- 11) *Kluyveromyces marxianus*-based platform for direct ethanol fermentation and recovery from cellulosic materials under air-ventilated conditions. Matsuzaki C., Nakagawa A., Koyanagi T., Tanaka K., Minami H., Tamaki H., Katayama T., Yamamoto K. and Kumagai H. *Journal of Bioscience and Bioengineering* 113, 604-607 (2012).
 - 12) A bacterial platform for fermentative production of plant alkaloids. Nakagawa A., Minami H., Kim JS., Koyanagi T., Katayama T., Sato F. and Kumagai H. *Nature Communications* 2, 326 (2011).
 - 13) Role of a PA14 domain in determining substrate specificity of a glycoside hydrolase family 3 β -glucosidase from *Kluyveromyces marxianus*. Yoshida E., Hidaka M., Fushinobu S., Koyanagi T., Minami H., Tamaki H., Kitaoka M., Katayama T. and Kumagai H. *Biochemical Journal* 431, 39-49 (2010).
 - 14) Purification, crystallization and preliminary X-ray analysis of β -glucosidase from *Kluyveromyces marxianus* NBRC1777. Yoshida E., Hidaka M., Fushinobu S., Koyanagi T., Minami H., Tamaki H., Kitaoka M., Katayama T. and Kumagai H. *Acta Crystallographica* F65, 1190-1192 (2009).
 - 15) Enantiomeric Separation of Racemic 1-Benzyl-*N*-methyltetrahydroisoquinolines On Chiral Columns and the Chiral Purity Determination of the *O*-Methylated Metabolites in Plant Cell Cultures by HPLC-CD On-Line Coupling in Combination with HPLC-MS. Iwasa K., Doi Y., Takahashi T., Cui W., Nishiyama Y., Tode C., Moriyasu M., Takeda K., Minami H., Ikezawa N. and Sato F. *Phytochemistry* 70, 198-206 (2009).
 - 16) Isolation of herbicide-resistant 4-hydroxyphenylpyruvate dioxygenase from cultured *Coptis japonica* cells. Liang Y., Minami H. and Sato F. *Bioscience, Biotechnology, and Biochemistry* 72, 3059-3062 (2008).
 - 17) Microbial production of plant benzylisoquinoline alkaloids. Minami H., Kim J.S., Ikezawa N., Takemura T., Katayama T., Kumagai H. and Sato F. *Proceedings of the National Academy of Sciences of the United States of America* 105, 7393-7398 (2008).
 - 18) A rapid, simple, and effective method for constructing a randomly mutagenized plasmid library, free from ligation. Koyanagi T., Yoshida E., Minami H., Katayama T. and Kumagai H. *Bioscience, Biotechnology, and Biochemistry* 72, 1134-1137 (2008).
 - 19) Functional analysis of norcochlorogenic acid synthase in *Coptis japonica*. Minami H., Dubouzet E., Iwasa K. and Sato F. *Journal of Biological Chemistry* 282, 6274-6282 (2007).
 - 20) γ -Glutamyltranspeptidase, but not YwrD, is important in utilization of extracellular glutathione as a sulfur source in *Bacillus subtilis*. Minami H., Suzuki H. and Kumagai H. *Journal of Bacteriology* 186, 1213-1214 (2003).
 - 21) Use of bacterial γ -glutamyltranspeptidase for enzymatic synthesis of γ -D-glutamyl

- compounds. Suzuki H., Izuka S., Minami H., Miyakawa N., Ishihara S. and Kumagai H. *Applied and Environmental Microbiology* 69, 6399-6404 (2003).
- 22) A mutant *Bacillus subtilis* γ -glutamyltranspeptidase specialized in hydrolysis activity. Minami H., Suzuki H. and Kumagai H. *FEMS Microbiology Letters* 224, 169-173 (2003).
- 23) Salt-tolerant γ -glutamyltranspeptidase from *Bacillus subtilis* 168 with glutaminase activity. Minami H., Suzuki H. and Kumagai H. *Enzyme and Microbial Technology* 32, 431-438 (2003).
- 24) Purification, cloning, and three-dimensional structure prediction of *Micrococcus luteus* FAD-containing tyramine oxidase. Roh J. H., Wouters J., Depiereux E., Yukawa H., Inui M., Minami H., Suzuki H. and Kumagai H. *Biochemical and Biophysical Research Communications* 268, 293-297 (2000).

総説

- 1) Fermentative production of plant benzylisoquinoline alkaloids in microbes. Minami H. *Bioscience, Biotechnology, and Biochemistry* 77, 1617-1622 (2013).
- 2) 「酵母で抗マラリア薬をつくる」、南博道、*化学* 68, 60-61 (2013).
- 3) 「大腸菌を用いた植物アルカロイドの発酵生産」、中川 明、南博道、*バイオサイエンスとインダストリー* vol.70 p.213-215、バイオインダストリー協会、(2012).
- 4) 「微生物による高等植物アルカロイドの生産」、南博道、佐藤 文彦、*BIOINDUSTRY* vol.29 p.53-59、シーエムシー出版、(2012).
- 5) 「微生物発酵の新展開」、南博道、*生物工学会誌* 第89巻・第7号 p.413、日本生物工学会、(2011).
- 6) 「微生物による高等植物アルカロイドの生産:ケシアルカロイドの微生物生産の可能性」、南博道、佐藤 文彦、*化学と生物* Vol.47 No.8, p.528-530、日本農芸化学会(学会出版センター)、(2009).
- 7) 「枯草菌の γ -グルタミルトランスぺプチダーゼに関する研究」、南博道、鈴木 秀之、熊谷 英彦、*応用微生物学研究* vol.1 p.109-114、応用微生物学研究協議会、(2003)

著書

- 1) Bioengineering of Isoquinoline Alkaloid Production in Microbial Systems. *New Light on Alkaloid Biosynthesis and Future Prospects* (Advances in botanical research). (Book chapter; vol.68) Nakagawa A., Matsumura E., Sato F. and Minami H. (2013) p.183-204 Academic Press.
- 2) Microbial production of plant benzylisoquinoline alkaloids. *Handbook of Natural Products* vol 1 Matsumura E., Matsuda M., Sato F. and Minami H. (2013) p.3-24 Springer.

- 3) Microbial production of plant benzyloisoquinoline alkaloids. *Handbook of Natural Products* vol 1 Matsumura E., Matsuda M., Sato F and Minami H. (2012) in press Springer.
- 4) Microbial Expression of Alkaloid Biosynthetic Enzymes for Characterization of Their Properties. Plant Secondary Metabolism Engineering: Methods and Applications (*Methods in Molecular Biology*). (Book chapter; vol.643) Minami H., Ikezawa N. and Sato F. (2010) p.111-120 Humana Press.
- 5) Effective use of bacterial γ -glutamyltranspeptidase to improve the taste of food. *State-of-the-art in flavour chemistry and biology* Suzuki H., Minami H., Izuka S., Miyakawa N., Kajimoto Y., Kato K., Ishihara S. and Kumagai H. (2004) p.205-212.